

KEEP OUT OF REACH OF CHILDREN

CARTAP-
HYDROCHLORIDE

GROUP

14

INSECTICIDE



ADABLAST

CARTAP

•INSECTICIDE•

FOR THE CONTROL OF INSECT PESTS OF RICE, CABBAGE, POTATO AND OTHER
CROPS SPECIFIED UNDER DIRECTIONS FOR USE

50SP



ACTIVE INGREDIENT:

Cartap hydrochloride -----500g/Kg

INERT INGREDIENTS:

Sodium dodecyl benzene sulfonate -----41.7g/Kg

White carbon black-----72.0g/Kg

Brilliant Blue FCF -----0.41g/Kg

Sodium sulfate -----to make 1000g

1Kg
Net Content

REGISTERED WITH THE FERTILIZER AND PESTICIDE
AUTHORITY PURSUANT TO P.D.1144
FPA REGISTRATION NO.:

Distributed by:
ADA MANUFACTURING CORPORATION
Rizal Avenue Extension, Brgy. Cabaruan,
Cauayan City, Isabela.



CAUTION





ADABLAST

CARTAP INSECTICIDE

WARNING: this product may be harmful if swallowed, inhaled or absorbed through the skin

PRECAUTIONS: Store in original container, tightly closed away from foods and feedstuffs. Do not smoke, eat or drinks while handling the product. Avoid inhalation of spray mist and skin and eye contact with the concentrate. Wear protective clothing and gloves while handling the product. Wash hands, arm and face thoroughly with soap and water after work. Avoid contamination of water supplies with chemical or container.

BABALA: Ang produktong ay Maaaring makalason kapag nalunok, nalanghap, o napunta sa balat. Huwag kumain, uminom o manigarilyo habang nagbobomba. Iwasang malanghap, malagyan ang mga mata at madikit sa balat ang kemikal. Magsuot ng damit panangga habang ginagamit ito. Hugasan ang sabon at tubig ang kamay, braso at mukha bago kumain o manigarilyo. Huwag gamitin uli ang pinagsidlan ng produkto. Iwasang maapektuhan ng kemikal at ng mga sisidlan nito ang anumang pinagkukunan ng tubig. Ang produktong ito ay nakakalason sa isda.

SIGN AND SYMPTOMS OF POISONING

WITH ALCOHOL INGESTION: Flushing, sweating, shortness of breath chest pain, hypotension.

WITHOUT ALCOHOL INGESTION: Dermatitis-itching and redness, shortness of breath, nausea, vomiting, muscle weakness paralysis.

FIRST AID TREATMENT:

IF SWALLOWED AND PATIENT IS CONSCIOUS, give 1 glass of water and induce vomiting by tickling back of throat or back of tongue with finger. Repeat procedure until vomit fluid is clear. Note: Never give anything by mouth to an unconscious person.

IN CASE OF SKIN CONTACT, remove contaminated clothing and wash affected areas with plain soap and water. If inhaled, remove from exposure area and bring patient to fresh air. If breathing has stopped, apply artificial respiration. If eyes are contaminated, Flush affected eyes with clean running water for at least 15 minutes. If symptoms persist, bring patient to the nearest medical clinic.

NOTE TO THE PHYSICIAN: The active ingredient of this product is a CNS ganglionic blocker. In case of severe exposure give intravenous injection of 100-200mg of L-cysteine or an intramuscular injection of 20-60 mg of BAL (dimercaprol or 2,3 dimercapto-1-propanol) Apply steroids against dermatitis.

MIXING: ADABlast Cartap 50 SP is generally recommended to be applied at the rate of ½ to 1kg product in 300-600 liters water per hectare. For specific pest and crop, consult the dilution guide above.

CROPS	PEST	TBSP PER 16 L WATER	APPLICATION
Cabbage/ Wongbok	Diamondback moth, cabbage moth, cabbage worm, Aphids	1.5-2.0	Spray as soon as pest damage becomes significant Repeat every 7 to 10 days Interval until pest damage is manageable
Cucurbits	Thrips, aphids, leafworm, flea beetle, squash beetle	1.5-2.0	
Onion/ Garlic	Thrips, cutworm	1.5-2.0	
Pechay	Looper, cutworm, aphids	1.5-3.0	
Potato	Thrips, aphids, cutworms, tuber moth, leafhopper, leaf miner	1.5-2.0	
Rice	Stemborer, green leafhopper, whorl maggots, brown planthopper	3.0-5.0	
Mango	Leafhopper	1.5-2.0	

1 tbsp = 10 gm

PRE-HARVEST INTERVAL: Do not spray within 10 days before harvest

RE-ENTRY PERIOD: Do not enter treated field within 2 days after spraying.

STORAGE AND DISPOSAL: Store in a cool dry place away from foodstuffs and animal feeds. Puncture and Triple rinse before returning to accredited hazardous waste collector. Do not re-use empty container.

PROHIBITION/WARRANTY STATEMENT

It is a violation of FPA rules and regulations to use this product in a manner inconsistent with its label. No warranty of any kind, expressed or implied, is made concerning the use of this product. User assumes all risks and liability resulting from improper handling use or application.

IN CASE OF EMERGENCY, PLEASE CONTACT NATIONAL POISON MANAGEMENT AND CONTROL CENTER- UP PGH, Manila

Emergency Tel No: (02)8524-1078; (02)8521-8450
Local 2311
24-hour Poison Information Service (02)8521-8451 loc.2311

VICENTE SOTTO MEMORIAL MEDICAL CENTER

B. Rodriguez St. Cebu City Tel (032)8253 9896

SOUTHERN PHILIPPINES MEDICAL CENTER (Formerly DAVAO MEDICAL CENTER)

JP Laurel Avo. Bajada Davao City Tel: (082)8227-2731

LOT/BATCH NO.:
DATE FORMULATED:
DATE OF EXPIRATION:

MANUFACTURED BY:
ZHEJIANG TONGLU HUIFENG
BIOSCIENCES CO., LTD
Hangzhou Zhejiang China