

KEEP OUT OF REACH OF CHILDREN



ADAMCO

70WP

ADACLEAR

•SNAIL KILLER•

FOR THE CONTROL OF "GOLDEN KUHOL" INFESTING RICE



Active ingredient:

Niclosamide ethanolamine salt -----700 g/kg
(equivalent to Niclosamide ethanolamine salt 83.4 g/kg)

Talc (Carrier) -----To make 1000g

35g
Net Content

REGISTERED WITH THE FERTILIZER AND PESTICIDE
AUTHORITY PURSUANT TO P.D. 1144

FPA Registration No.:

Lot/Batch No.

Mfg Date:

Expiry Date:

Distributed by:
ADA MANUFACTURING CORPORATION
Rizal Avenue Extension, Brgy. Cabaruan,
Cauayan City, Isabela.



ADACLEAR

ADAMCO

SNAIL KILLER FOR THE CONTROL OF "GOLDEN KUHOL" INFESTING RICE

DIRECTIONS FOR USE:

Level field properly. Irrigate rice field to allow snails to surface out and maintain 3 cm depth of water.

Spray **ADAClear 70 WP** at 350 grams per hectare. Do not irrigate further for at least 48 hours. Follow standard irrigation practice after this treatment.

CROP	PEST	Grams per tank load	No. of Tank Load per Ha
RICE	Golden Apple Snail	35 grams (1 sachet)	10

1 tbsp = 10 grams 1 Tank Load = 16 Liters

Frequency & Method of Application:

Spray **ADAClear 70 WP** during high level of infestation between 0-15 days after seeding/ days after transplanting.

MIXING: Pour the required amount of **ADAClear 70 WP** in a sprayer tank partially filled with water and stir. Fill the tank with the required amount of water and mix well.

Re-entry Period: 48 hours after spraying.

Compatibility: Do not mix **ADAClear 70 WP** with other products.

STORAGE AND DISPOSAL: Do not store near seeds, food and other feedstuff. Do not re-use container. Triple rinse the empty container before disposal.

PROHIBITION AND WARRANTY STATEMENT: It is a violation of FPA rules and regulations to use this product in a manner inconsistent with its label. No warranty of any kind, expressed or implied, is made concerning the use of this product. User assumes all risks and liabilities resulting from improper handling, use or application.

WARNING: This product may be harmful if swallowed, inhaled or absorbed through the skin.

PRECAUTIONS:

Store tightly in its closed container. Keep away from foods and feedstuffs. Do not allow the product to come into contact with skin and wear waterproof protective clothing including boots and gloves. After spraying, wash hands and contaminated parts of the body thoroughly with plenty of soap and water. After use, clean all spray equipment thoroughly with water. Bury empty containers. Do not allow the chemical or container to contaminate sources of water. This product is toxic to fish.

BABALA: Iwasan na makapasok sa mga mata ang alikabok o tilamsik nito. Kung madikit sa balat, hugasan ng maraming tubig at sabon ang lahat ng parte ng katawan na naapektuhan ng kemikal. Kung makapasok sa mata padaanan ng maraming tubig sa loob ng 15 minuto at dalhin sa doktor. Maghugas ng kamay bago kumain at pagkatapos ng trabaho. Hugasang maigi ang lahat ng kasangkapang ginamit sa pagbobomba. Huwag hayaang maapektuhan ng kemikal o ng sisidlan nito ang anumang pinagkukunan ng tubig. Ang produktong ito ay nakakalason sa isda.

SIGNS AND SYMPTOMS OF POISONING:

Ataxis or uncoordinated movements, staggering, dyspnea and vomiting. If eye contact occurs, the conjunctive will redden and swell.

FIRST AID TREATMENT:

If on Skin: Remove contaminated clothing and wash affected areas thoroughly with plain soap and water.

If in Eyes: Hold eyelids open and flush eyes with clean running water for at least 15 minutes.

If Inhaled: Remove from contaminated area and bring to fresh air. If breathing has stopped, apply artificial respiration.

If Swallowed: Give 1-2 glasses of water to drink. Induce vomiting by tickling the back of the throat or pressing back of the tongue with a finger. Repeat procedure until the vomit fluid is clear. Never give anything by mouth to an unconscious person.

GET MEDICAL ATTENTION IMMEDIATELY

NOTE TO THE PHYSICIAN:

There is no specific antidote. If illness results from exposure, treat symptomatically. If ingested orally, give 200 ml liquid paraffin and gastric lavage with 4 liters of water. Afterwards, treat with activated charcoal and sodium sulfate. In case of mucous membrane reactions, treat symptomatically with antihistamine gel and corticosteroid formulations

Manufactured by:

Zhejiang Tonglu Huifeng Biosciences Co., Ltd
Hangzhou Zhejiang China

IN CASE OF EMERGENCY PLEASE CONTACT:

Davao Medical Center: Bajada, Davao City
Tel.No.: (082)8227-2731

National Poison and Information Service: PGH,
Manila Tel. Nos.: (02) 8524-1078; (02) 8521-8450 loc.2311