

**KEEP OUT OF REACH OF CHILDREN**

GLYPHOSATE

GROUP

**G**

HERBICIDE



# ADABLAST GLYPHO

•HERBICIDE•

**480SL**



**ACTIVE INGREDIENT:**

Glyphosate IPA----- 480 g/L

**INERT INGREDIENTS:**

Alkyl Polyglucoside----- 95.82 g/L

Ethoxylated tallow amine----- 48.60 g/L

Water (Solvent)----- to make 1L

**1L**  
Net Content

REGISTERED WITH THE FERTILIZER  
AND PESTICIDE AUTHORITY  
PURSUANT TO P.D. 1144

FPA Registration No.:

Distributed by:  
**ADA MANUFACTURING CORPORATION**  
Rizal Avenue Extension, Brgy. Cabaruan,  
Cauayan City, Isabela.



# ADABLAST GLYPHO

## READ ENTIRE LABEL BEFORE USING THE PRODUCT

### A. CROPPING SYSTEMS

\* Blanket spraying prior to planting vegetables, corn rubber, oil palm, coconut, mango, pineapple, citrus, coffee, cacao, banana and other perennial tree crops.

\* Minimum or reduced tillage systems for sugarcane, rice, corn and other cereal and vegetables crops.

\* Blanket spraying, strip spraying, spot spraying or wiping around young and mature rubber, oil palm, coffee, sugarcane, citrus, mango, cacao, coconut and banana.

### B. NON-CROP USES

Reforestation areas, subdivision lots, rice levees, drainage ditches, irrigation canals, around fish ponds, field boundaries, roadsides, fence rows, railroad yards, railroad tracks, around buildings, golf courses and memorial parks.

**WARNING:** This product may be harmful if swallowed, inhaled or absorbed through the skin.

**SAFETY PRECAUTIONS:** Store in original container, tightly closed away from foods and feedstuffs. Do not smoke, eat or drink while handling the product. Avoid inhalation of spray mist and skin and eye contact with the product and diluted material. Wear eye goggles, gloves and overalls when handling the undiluted product. After work, thoroughly wash hands, face and contaminated skin using plenty of soap and water. Avoid contamination of water supplies with chemical or container.

**BABALA:** Ang produktong ito ay Maaaring makasama sa kalusugan kapag nalunok, nalanghap, o nadampi sa balat. Ilagay sa orihinal na lalagyan malayo sa pagkain at ito ay mahigpit na takpan. Huwag manigarilyo, kumain o uminom at magsuot ng damit pananggalang habang ginagamit ang produkto. Iwasang malanghap o malagyan ng kemikal ang balat at mata. Hugas ng mabuti ang mga kamay, mukha at lahat ng parte ng katawan matapos magbomba. Iwasang maapektuhan ng kemikal at lalagyan nito ang anumang daluyan o pinagkukunan ng tubig.

**SIGNS AND SYMPTOMS OF POISONING:** Symptoms of the acute poisoning such as weaknesses, sweating, nausea, vomiting and stomach ache were visible right after the working period.

**FIRST AID TREATMENT:** If swallowed, dilute by drinking water or milk and call a physician. If on skin, remove contaminated clothing and wash affected areas with plain soap and water. If on eyes, immediately flush with running water for at least 15 minutes.

**NOTE TO PHYSICIAN:** Symptomatic and supportive treatment.

### IN CASE OF EMERGENCY, CONTACT:

**NATIONAL POISON MANAGEMENT AND CONTROL CENTER c/o UP-PGH.**  
Tel. Nos. +632 8524-1078/8567-2057, +632 8521-8451 loc. 2311

**VICENTE SOTTO MEMORIAL MEDICAL CENTER**  
B. Rodriguez St. Cebu City, Tel. (032) 8253-9891 to 99

**DAVAO MEDICAL CENTER, Bajada, Davao City**  
Tel. No.: (082) 8227-2731, (082)8226-4017

### DIRECTIONS FOR USE:

Add the required amount of **ADABlast Glypho 480 SL** with the 1/2 required amount of water in the mixing spray tank. Mix well before adding the remaining portion of water. For knapsack sprayers, it is suggested to mix the solution in a separate container and fill the knapsack sprayer with the mixed solution.

In early dry season, low volume spraying of **ADABlast Glypho 480 SL** applied at 200-400 liters of water per hectare gives best results. When sprayed uniformly to tall, dense foliage of cogon or on shorter species such as mutha, **ADABlast Glypho 480 SL** will provide a minimum of 7 months weed control.

CROPS REGISTERED	WEEDS CONTROLLED	LITERS PER HECTARE	RATE OF ADABlast Glypho APPLICATION VOLUME (Tbsp*/16 L SPRAYER)	
			12.5 TANK LOADS OR 200 L/HA	25 TANK LOADS OR 400 L/HA
GLYPHOSATE TOLERANT CORN <i>Non-crop &amp; minimum or reduced tillage</i>	Itch grass, Morning Glory, Dayflower, Syndrella, Button weed, Spindle Top, Calopogonium, Harry Beggar ticks, Tropic Ageratum	2-3	16-24	8-12
MANGO, BANANA, COFFEE, CITRUS	COGON	3.0	24	12
	Shade Conditions	4.5	36	18
RUBBER, OIL PALM, COCONUT, CITRUS	MUTHA	2.0	16	8
	PARAGRASS	2.0	16	8
	TALAHIB	4.5	36	18
	FERN	2.0	16	8
SUGARCANE AND VEGETABLES	Shade Conditions	2.0	16	8
	OTHER ANNUAL WEEDS	2.0	16	8

**1 tbsp ADABlast Glypho 480 SL = 10 mL**

In mixed weeds situations, 2 liters of **ADABlast Glypho 480 SL** per 200 liters water sprayed evenly on weeds will provide effective control.

A 3-30% aqueous solution of **ADABlast Glypho 480 SL** may be applied with wiper to control weeds around sensitive plants like young coffee, cacao, citrus, mango, banana, sugarcane and vegetables. For Glyphosate-tolerant corn varieties, apply **ADABlast Glypho 480 SL** at 20 to 25 Days After Seeding (DAS). And at 40-45 DAS, at half the above rate per application timing. Use the higher recommended rate when weed population is high and weed height exceeds beyond knee high.

**COMPATIBILITY/INCOMPATIBILITY:** Incompatibility with paraquat and diquat

**STORAGE AND DISPOSAL:** Do not store near seeds, food and other feedstuff. Do not re-use container. Triple rinse the empty container before disposal.

**PROHIBITION AND WARRANTY STATEMENT:** It is a violation of FPA rules and regulation to use this product in a manner inconsistent with its label. No warranty of any kind, expressed or implied, is made concerning the use of this product. User assumes all risks and liabilities resulting from improper handling, use or application.

Date of Formulation:  
Date of Expiration:  
Lot/Batch No.:

MANUFACTURED BY:  
ZHEJIANG TONGLU  
HUIFENG BIOSCIENCES CO., LTD  
Hangzhou Zhejiang China