

WARNING: THIS PRODUCT IS HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN. The product may cause moderate skin irritation and mild eye irritation.

PRECAUTIONS: Wear overalls or long-sleeved shirt, long trousers and gloves when mixing; wear overalls or long-sleeved shirt and long trousers and solid footwear when spraying. Avoid breathing in spray mist and accidental skin contact. Immediately wash off accidental splashes of the product or spray mixture. After use, wash entire body and change clothes. Do not eat, drink or smoke while handling the product; wash hands before doing so. Wash protective equipment before re-use. Store in original container, tightly closed, away from food and feeds.

BABALA: ang produktong ito ay nakakalason kapag nalunok, nalanghap o nadikit sa balat. Iwasang mabilad sa araw o mahamugan. Itabi sa maaliwalas na lugar at ilayo sa pagkain o patuka. Iwasang manitan ng higit sa 30-degrees celcius. Mag ingat sa paghalo at paggamit ng produkto upang maiwasang malagyan ang mga mata, balat o damit. Magsuot ng damit pananggalang habang humahawak ng produkto. Maligo at magpalit ng damit pagkatapos magtrabaho. Labhang hiwalay ang damit pananggalang. Huwag kumain, uminom o manigarilyo habang gumagamit ng produkto. Maghugas ng kamay at mukha bago kumain, uminom o manigarilyo.

SIGNS AND SYMPTOMS OF POISONING:

For lambda-cyhalothrin, ingestion may produce nausea, vomiting, abdominal pain and diarrhea. Severe cases may result to ataxia, convulsion and coma. May also cause skin and eye irritation. Exposure of the face to the product concentrate may cause a transient tingling or burning sensation. For thiamethoxam, no case of human poisoning is on record. Under laboratory conditions, unspecific poisoning symptoms were observed in rats.

FIRST AID: If poisoning is suspected, call a physician immediately. After accidental skin contact strip of contaminated clothing and wash affected area with plenty of water and soap. Rinse splashes from eyes with abundant fresh and clean water for at least 15 minutes. If inhaled, move to clean air. If breathing is irregular or stopped, administer artificial respiration. "If ingested, seek medical advice immediately and show this container or label. Do not induce vomiting."

NOTE TO PHYSICIAN: If ingested, carry out gastric lavage with care to prevent aspiration. Administer symptomatic and supportive treatment.

KEEP OUT OF REACH OF CHILDREN

LAMBDA-CYHALOTHRIN + THIAMETHOXAM	GROUP	3A+4A	INSECTICIDE
-----------------------------------	-------	-------	-------------



ADABLAST LAMBDA + TMX

INSECTICIDE

Insecticide for the control of sucking and chewing pests in potato, string beans, eggplant, ampalaya, corn, mango and banana and rice.

247 ZC

250ml
Net Content



Active Ingredient:	
Lambda-cyhalothrin	106 g/L
Thiamethoxam	141 g/L
Inert Ingredients:	
Propylene glycol	45.7 g/L
NNO	18.2 g/L
Calcium Lignosulfonate	17.4 g/L
Xanthan Gum	1.7 g/L
Polydimethylsiloxane	1.6 g/L
Water	To make 1 Liter

REGISTERED WITH THE FERTILIZER AND PESTICIDE AUTHORITY PURSUANT TO P.D. 1144
FPA Registration No.:

IN CASE OF EMERGENCY, PLEASE CONTACT: NATIONAL POISON MANAGEMENT AND CONTROL CENTER

UP PGH, Manila Emergency Tel Nos:
(02)8524-1078. (02)8554-8400 Loc. 2311

24-hour Poison Information Service
(02)8521-8451 Loc. 2311

VICENTE SOTTO MEMORIAL MEDICAL CENTER
B. Rodriguez St. Cebu City Tel: (032)8253-9898

BEFORE APPLYING THE PRODUCT, READ AND UNDERSTAND THE LABEL"

DIRECTIONS FOR USE: ADABlast Lambda+TMX
247 ZC is registered for use on crops and pests listed below. Mix with water as indicated. For vegetables and corn, apply using a 16-liter knapsack sprayer using a spray volume of 480 liters/ha or 30 tank loads per hectare. For mango, apply using a power sprayer. For banana, apply as bud injection at an average volume of 60 ml solution/bd. For rice, apply using a 16-liter knapsack sprayer at spray volume of 192 liters/ha

CROPS	PEST	Recommended Rate ml product/ha	ml product/16L	Pre-Harvest Interval (PHI)
POTATO	Whiteflies, Aphids, Thrips	100-200	4.0 - 6.0	7 days
STRING BEANS	GLH, Aphids, Mites	100-200	4.0 - 6.0	90 days after flower induction
EGGPLANT	Green Leafhopper, Whiteflies Aphids	100-200	4.0 - 6.0	7 days
BITTER GOURD (Ampalaya)	Leafhopper, Aphids, Thrips, Fruit fly, Cutworm	100-400	4.0 - 13.0	3 days
CORN	Cutworm	100 to 400	10.0 - 16.0	3 days
MANGO	Mango Leafhoppers	20 - 30ml/ 100	N/A	3 days
BANANA	Flower Thrips	1-3 mL/L	N/A	90 days
RICE	Stemborer, Leafhoppers	100-200	10.0 - 30.0	14 days

1 tbsp - 10 ml

STORAGE AND DISPOSAL: Store in a cool place. Keep container tightly closed in a dry and well-ventilated place. Triple or preferable pressure rinse containers before disposal and deliver empty pack-aging to an approved waste management facility.

PROHIBITION AND WARRANTY: It is a violation of FPA rules and regulations to use this product in a manner inconsistent with its label. No warranty of any kind, expressed or implied, concerning the use of this product. User assumes all risks and liabilities resulting from handling, use or application.

Date of Formulation:
Date of Expiration:
Batch No.:

Manufactured by:
Zhejiang Tonglu Hui Feng Biosciences Co., LTD
Hangzhou Zhejiang China