

KEEP OUT OF REACH OF CHILDREN

**COPPER
HYDROXIDE**

GROUP

MO1

FUNGICIDE



ADACOP
770WP

• FUNGICIDE •



Active Ingredient:
Copper hydroxide-----770 g/Kg

Inert Ingredients:
Sodium poly-----9.08 g/Kg
NNO-----4.40 g/Kg
Kaolin-----To make 1 Kg

500g
Net Content

REGISTERED WITH THE FERTILIZER
AND PESTICIDE AUTHORITY
PURSUANT TO P.D. 1144

FPA Registration No.:

Distributed by:
ADA MANUFACTURING CORPORATION
Rizal Avenue Extension, Brgy. Cabaruan,
Cauayan City, Isabela.

CAUTION



ADACOP

FUNGICIDE

WARNING: This product maybe harmful if swallowed, inhaled or absorbed through the skin. May cause skin and eye irritation.

PRECAUTIONS: May cause irritation of eyes, nose, throat and skin. May be harmful if inhaled or swallowed. Avoid breathing of dust and spray mist. Avoid contact with skin, eyes and clothing. In case of contact, wash skin or eyes with plenty of water.

BABALA: Maaaring pagmulan ng panghahapdi ng mga mata, ilong, lalamunan at balat. Maaaring makasama kapag nalanghap o nalunok. Iwasang makalanghap ng alikabok o tilamsik nito. Kung sakaling malagyan nito, hugasang mabuti ang balat at mga mata.

SIGNS AND SYMPTOMS OF POISONING: Vomiting, colic, diarrhea and dehydration.

FIRST AID TREATMENT:

IF SWALLOWED: Call a physician. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything to an unconscious person.

IF ON SKIN: Wash with plenty of soap and water. If irritation persists, get medical attention.

IF INHALED: Bring victim to fresh air. If not breathing, apply artificial respiration, preferably mouth-to-mouth. Get medical attention.

IF IN EYES: Flush with plenty of water for at least 15 minutes. Get medical attention.

NOTE TO PHYSICIAN: Give supportive and symptomatic therapy.

IN CASE OF EMERGENCY, PLEASE CONTACT

National Poison Control and Information Service
PGH, Manila
Tel. Nos.: (02)8524-1078, (02)8554-8400 loc. 2311
c/o the UP-Philippine General Hospital

PROHIBITION AND WARRANTY STATEMENT

"It is a violation of the FPA Rules and regulations to use this product in a manner inconsistent with its label. No warranty of any kind, expressed or implied is made concerning the use of this product. User assumes all risks and liability resulting from handling, use or application.

DIRECTION FOR USE

Apply ADACop 770 WP at the following recommended rates per 16 liters of water. Mix the required amount of ADACop 770 WP into a little water to make a slurry and then pour the concentrated suspension into a measured quantity of water in the spray tank and stir vigorously. Spray at 7-to-14-day intervals depending on disease severity.

CROPS	DISEASES	RECOMMENDED RATES (Tbsp/ 16L)
Rice	Rice Blast, Sheath Blight, Bacterial Leaf Streak	2.0 – 5.0
Mango	Anthrachnose	2.0 – 6.0
Cacao	Black Pod	2.0 – 6.0
Coffee	Leaf Rust, Coffee Berry Disease	2.0 – 6.0
Citrus	Melanose, Scab, Canker, Fruit Rot, Brown and Black Spot	2.0 – 6.0
Tomatoes	Early Blight, Fruit Rot, Downy Mildew, Septoria	2.0 – 6.0
Cabbage	Black Rot, Black Leaf Streak	2.0 – 6.0
Carrots	Carrot Blight, Alternaria	2.0 – 6.0
Onions	Purple Blotch, Downy Mildew	2.0 – 6.0
Beans	Rust, Bacterial Blight	2.0 – 6.0
Banana	Sigatoka	2.0 – 6.0

1 Tbsp = 10 grams

PRE- HARVEST INTERVAL: Allow 7 days from last spray till harvest.

RE-ENTRY PERIOD: Do not enter treated area until spray has dried.

COMPATIBILITY: ADACop 770 WP is compatible with most insecticides and fungicides. However, it should not be used with synthetic pyrethroid based insecticides.

STORAGE AND DISPOSAL: Do not store near seeds, food and other feedstuff. Do not re-use container. Puncture and triple rinse before returning to accredited hazardous waste collector. Avoid contamination of water supply with chemical or empty container.

FORMULATION DATE:
EXPIRATION DATE
LOT/BATCH NO.:

Manufactured by:
Sales y Derivados de Cobre S.A. (SALDECO)
Calle 4, Mz-B1, Lote 18, Urb.
Industrial Las Vegas,
Puente Piedra Lima 22 (Peru)

